

Illustrated identification key to the bats of Europe

By Christian Dietz
& Otto von Helversen

Electronic publication

Version 1.0

First released 15.12.2004

© Dietz & von Helversen

Illustrated identification key to the bats of Europe

By Christian Dietz & Otto von Helversen

Electronic Publication

Version 1.0. released 15.12.2004

Tuebingen & Erlangen (Germany)

with 228 photographs by Christian Dietz

and Otto von Helversen

and 14 drawings by Otto von Helversen.

Copyright © 2004 by the authors

Christian Dietz & Otto von Helversen.

All rights reserved.

The text, or parts thereof, and the photographs
must not be reproduced without written permission
of the authors.

Authors' email addresses:

ChristianDietzHorb@web.de

Helver@biologie.uni-erlangen.de

Illustrated identification key to the bats of Europe

By Christian Dietz & Otto von Helversen

PREFACE

Since the identification key on the European bats published by VON HELVERSEN in 1989 a surprising number of new species has been discovered in Europe¹⁾ with the number of species rising from 31 to 39 in 2004. This rapid development in taxonomy and systematics has made it harder for field biologists to identify living bats, especially in the Mediterranean area. Most of the newly discovered cryptic species are closely related to one or more long known species. In some of these species groups identification has been problematic for many years and species assignment could only be solved by the aid of modern molecular methods. But the analysis of genetic characters is an inappropriate method for most field studies. Our new key on the European bats was written mainly for biologists aiming to identify captured living bats which would be released after identification.

The first part of the identification key is written mainly for students and beginners in the studies of bats, as most of the groups can be identified quite easily. The separated keys to the more difficult groups are addressed mainly to the more experienced field workers to help them when faced with an unknown species or with the most difficult groups of very similar bats.

However, not all characters of the newly described species are currently known in their full variability and furthermore some taxonomic questions are not finally clarified.

¹ We define Europe as the continental area west of a line from the Crimea peninsula to the north and the Mediterranean islands of Crete, Malta, Sicily, Sardinia, Corsica and the Balearic Islands.

Therefore we have chosen an electronic publication which is easier to update with new characters or newly described species. This will ensure that the identification key remains up to date. We would like to encourage all bat workers using our key to tell us about their experiences and to pass on their comments and thus help us keep the key up to date.

ACKNOWLEDGEMENTS

Many people helped us by sharing their experience in the field identification of bats. We are grateful to all our friends and colleagues for the many discussions on their newest findings in taxonomy and genetics or for checking characters in various parts of Europe.

We also benefited greatly from many people who helped us to capture the bat species so that we could take measurements and photographs. We are specifically grateful to Mauro Mucedda, Maja Zgmajster & Alenka Petrinjak for their help in catching *Plecotus sardus*, *Myotis punicus* and *Plecotus macrobullaris*. Isabel Schunger joined us during most of our excursions and gave valuable help in all respects. Further help to get permissions and to find and capture bats was provided by (in alphabetic order) Monika Braun, Kamen Christov, Philipp Dietz, Klaus Echle, Antoaneta Gueorguieva, Teodora Ivanova, Ingrid Kaipf, Vassiliki Kati, Spartak Merdschanov, Dessislava Merdschanova, Alfred Nagel, Rainer Nagel, Dietmar Nill, Eleni Papadatou, Boyan Petrov, Torsten Pröhl, Paul Schuhmacher and Nikolaj Simov. For many discussions on characters in identifying bats we are much obliged to Ursel Häussler, Ahmet Karataş and Katerina Tsytsulina. For their permanent help in identifying bats by molecular genetic methods and for extensive discussions we are grateful to Frieder Mayer and Andreas Kiefer. We are also indebted to Doris Mörike for giving us access to the collections of the Staatliches Museum für Naturkunde Stuttgart (SMNS).

We are particularly indebted to Paul Bates for proof-reading the manuscript and for the many corrections of the language. We thank him for many useful suggestions how to improve the descriptions and how to make them easier to understand.

HOW TO PROCESS A CAPTURED BAT

As this key is written to determine living bats in the hand it is necessary to mention first that bats are protected in all European countries. Therefore a licence is required to catch and handle bats.

Bats might be caught by a variety of techniques both at roost sites and in free flight. General advice in bat work and how to catch bats is given for example in the “Bat Workers' Manual” published by the Joint Nature Conservation Committee, also available for free in electronic format (www.jncc.gov.uk/Publications/bat_workers). Once bats are captured great care is needed to ensure that they are determined and measured quickly and without causing any harm. Pregnant or lactating females with attached young should be released immediately without further disturbance.

After being caught, bats can be best kept in soft cloth bags. Bags should be always hung up and never laid on the ground. Horseshoe bats and sexually active males of the large vespertilionid bats should always be kept as singles. For horseshoe bats the bags should be fixed in a way that allows the bats to hang head down and they should be kept captive as briefly as possible. Small vespertilionid bat species like pipistrelles or Daubenton's bats can be kept in small groups in bags, but species should never be mixed.

To obtain the bat's measurements and to examine the characters it is best to wrap them in a cloth or to hold them with soft gloves. Make sure you do not handle them too long, avoid holding a bat tight in your palm (if they are very active, they might suffer from heat stress). Never hold the bats by their forearms, elbows or wing tips only, since their flight muscles might be strained or, even worse, their skeletal system damaged.

WHICH ARE THE MEASUREMENTS USED IN THIS IDENTIFICATION KEY?

The main measurements (see table 1) are the lengths of forearm (FA), fifth finger (D5) and third finger (D3). Additional valuable measurements are the lengths of thumb (D1), lower leg (Tib), hind foot (HF). For some species groups, length and width of the ear and of the tragus are used (ear length (earL), ear width (earW), tragus length (tragL) and tragus width (tragW)). In other species groups, the length of some of the phalanges: 1st and 2nd phalanx of the 4th finger (P4.1 and P4.2) and the 2nd and 3rd phalanx of the 3rd finger (P3.2 and P3.3) are needed. In some very difficult species groups also the upper tooth row length (CM³) can be of some help.

Although measurements like wingspan, head-body-length and tail length are often mentioned in books, they are not really useful and there is too much variation through different measuring techniques, so they should be avoided to reduce unnecessary stress for the bats. Body mass is a good indicator for the identification of some species when taken at the same time of the day. However, it is omitted here since there are considerable changes in the course of a day and a year.

All measurements given in this key are only valid for fully grown (adult) individuals. At the time of their first flight, the bones of juveniles are not fully ossified. In not fully grown bats, the epiphyses are best visible in the joints of the digits against a light background. Small juveniles have long stretched joints and the fingers are still cartilaginous. With the onset of flight, most parts of the fingers are fully ossified, but the growth plates near the joints are apparent as a light (translucent) cartilaginous gap. In autumn the cartilage is replaced by bone and the joint becomes more and more rounded, knuckle-like (see Fig. 1 – 2, Fig. 1 shows an eight week old juvenile, Fig. 2 the same bat at the age of one year). In addition juveniles of most species are more greyish in coloration and often have a sparser fur (Fig. 3 – 4).

Table 1: Measurements used in the identification key.

measurement	abbreviation	best taken by	measurement used in
Forearm-length	FA	Caliper or steel ruler	All bats
Length of fifth finger	D5	Caliper or steel ruler	All bats
Length of third finger	D3	Caliper or steel ruler	All bats
Length of thumb	D1	Caliper or steel ruler	Whiskered bats, <i>Plecotus</i>
Length of tibia	Tib	Caliper or steel ruler	Whiskered bats, <i>Plecotus</i>
Length of hind foot	HF	Caliper or steel ruler	Whiskered bats, <i>Plecotus</i>
Length of ear	earL	Steel ruler	Large <i>Myotis</i>
Width of ear	earW	Steel ruler	Large <i>Myotis</i>
Length of tragus	tragL	Steel ruler	<i>Plecotus</i>
Width of tragus	tragW	Steel ruler	<i>Plecotus</i>
Length of 2nd phalanx of 3rd finger	P3.2	Caliper	<i>Pipistrellus pipistrellus</i> / <i>pygmaeus</i>
Length of 3rd phalanx of 3rd finger	P3.3	Caliper	<i>Pipistrellus pipistrellus</i> / <i>pygmaeus</i>
Length of 1st phalanx of 4th finger	P4.1	Caliper	Medium sized horseshoe bats
Length of 2nd phalanx of 4th finger	P4.2	Caliper	Medium sized horseshoe bats
Length of upper tooth row	CM ³	Caliper	Large <i>Myotis</i> , <i>Eptesicus serotinus</i> / <i>bottae</i> , <i>Plecotus austriacus</i> / <i>kolombatovici</i>



Plate 1: Age classification in bats. Juveniles are on the left, adults on the right.
M. aurascens (1 – 2), *M. emarginatus* (3), *R. ferrumequinum* (4).

HOW TO TAKE THE MEASUREMENTS USED FOR IDENTIFICATION

Measurements are of any value only when taken in the same standardised way. Calipers and in some measurements steel rulers will be needed to obtain reliable values.

To take the wing measurements (Fig. 8) it is best to hold the bat (for right-handed people) in your left palm curling your fingers around the bat's body (as shown in Fig. 5).

To take the **forearm length** (see Fig. 5) it might be easiest to keep the bat in your palm and to fix the folded right forearm of the bat with your thumb and the tip of your index finger. The inner end of the caliper can be fixed by a finger at the bat's elbow. The maximum forearm length is taken between the elbow and the wrist [this is the maximum forearm length (FA+), in some publications the forearm length is given without the wrist (FA-) representing the true length of the forearm bone. Usually the values of FA- are about 0.5 to 1.2 mm less than FA+, depending on the species. As it is much harder to reproduce reliable FA- measurements, we recommend to use only FA+ in future or to give both values]. It is important to ensure that the moveable jaws of the caliper are well attached to elbow and wrist and that the elbow is held parallel to the caliper.

To take the **lengths of the third and the fifth digit** it is easiest to keep the bat (for right-handed people) with your left hand and attach it, the bat's ventral side up, to a flat surface (table or one's thigh) and open the wing (Fig. 6 – 7). The outer end of the caliper is best attached to the inside of the wrist and the length to the tip of the finger is taken. In the fifth digit length the full length of the straight finger is taken, in the third finger length in living bats it is better to take the secant of the finger of the outstretched wing. The lengths of the phalanges are taken as shown in Fig. 11.

Thumb length is measured as the maximum distance of the straight thumb without claws (Fig. 9). **Hind foot length** is taken from the base of the spur to the toes without claws (Fig. 10). The **tibia length** is taken from the knee to the end of the tibia after having bent the foot (Fig. 10). **Ear width** in large *Myotis* is taken as shown in Fig. 12 and Fig. 13 as the combined value of a and b at the height of the tip of the tragus. **Tragus width** in *Plecotus* is taken at the point of the tragus with the maximum width (Fig. 14). The tragus is usually not flat, to obtain reliable values it is useful to attach the tragus to a steel ruler in order to make it level. **Tragus length** in *Plecotus* is measured from the notch at the outside of the tragus above its basal lobe to the tip of the tragus (Fig. 14). The **upper tooth row length** can also be measured in living bats, but experience and concentration are necessary not to hurt the bat. This measurement is only necessary in some species groups if the identification is not clear having used all other characters given in the key. It might be helpful to obtain these data in the species groups of *Myotis myotis* / *punicus* / *blythii* in some Mediterranean areas, *Plecotus austriacus* / *kolombatovici* along the Adriatic coast and Greece and in *Eptesicus serotinus* / *bottae* along the coastlines of Turkey and the Greek islands. This measurement is taken as the distance between the posterior margin of the last molar and the base of the canine (Fig. 15).

Plate 2: How to hold a bat to take measurements of forearm and fingers.

E. nilssonii (5 - 7).

Plate 3: How to take measurements of the wing, thumb and leg.

E. nilssonii (8), *P. auritus* (9), *M. capaccinii* (10), *P. pygmaeus* (11).

Plate 4: How to take measurements of the ear and the upper tooth row.

M. punicus (12), *M. blythii* (13), *P. kolombatovici* (14), *M. blythii* (15).












Plate 2: How to hold a bat to take measurements of forearm and fingers.





LIMITS OF SPECIES IDENTIFICATION

Unfortunately it is not always possible to determine all bats by external characters. Even when considering all characters given in this key some species are difficult to distinguish and even more, some individuals differ so much from the usual appearance that they do not match the given descriptions. There is a high degree of intraspecific variability within some species (and perhaps this identification key is not reflecting the whole range of variability).

In some groups taxonomic questions have not been solved yet. Some cryptic species may still be awaiting discovery, and vagrants or accidentally transported individuals might further increase the list of European bat species.

IDENTIFICATION KEY TO THE FAMILIES

1) tail extending up to a half beyond the narrow tail membrane (Fig. 18). Lower part of the posterior margin of the ear with pronounced lobes (Fig. 17). No nasal process or nose leaf (Fig. 16). Only one species in Europe. – [Molossidae](#)

► Tail included completely in the broad tail membrane or except for the last one or two vertebrae (maximum about 5 mm) (Fig. 21 and 24). – **2**

2) Nose with a pronounced nose leaf (cutaneous process) (Fig. 19). Ears without a tragus (Fig. 20). Tail shorter or of same length as the hind legs (Fig. 21). Echolocation calls audible by a bat detector as long whistles. Five species in Europe. – [Rhinolophidae](#)

► No nose leaf (Fig. 22). Ears with a tragus (Fig. 23 and 26). Tail longer than the hind legs (Fig. 24). – **3**

3) Ears projecting beyond the top of the head (Fig. 23). Second phalanx of the third finger not specially elongated (up to about twice the length of the first phalanx, usually both are of more or less the same length). 32 Species in 9 genera in Europe. – [Vespertilionidae](#).


► Ears very short and triangular, not projecting beyond the top of the head (as if cut with scissors) (Fig. 25 and 26). Second phalanx of the third finger (P3.2) about three times as long as the first phalanx (P3.1) (Fig. 27). 3rd and 4th fingers at rest folded in the joint between 1st and 2nd phalanges. Only one species in Europe. – [Miniopteridae](#).

Plate 5: The four European families of bats.

T. teniotis (16, 17, 18), *R. mehelyi* (19), *R. ferrumequinum* (20, 21),
E. serotinus (22), *M. aurascens* (23), *M. daubentonii* (24),
M. schreibersii (25, 26, 27).



Family Molossidae



Family Rhinolophidae



Family Vespertilionidae



Family Miniopteridae

A) RHINOLOPHIDAE

1) connecting process (= upper saddle process, Crista) bluntly rounded in profile and shorter than the lower tip of the sella (= lower saddle process) (Fig. 29 and 33). – **2**

▶ connecting process in profile pointed and always longer than the tip of the sella (Fig. 36, 40 and 44). – **3**

2) Biggest of the five horseshoe bats, FA > 54 mm (54.0 – 62.4 mm, lowest extreme 51.0 mm), D5: 63 – 77 mm, D3: 78 – 94 mm, P4.1: 9.5 – 13.4 mm; P4.2: 17.5 – 22.5 mm. High and bluntly rounded connecting process, the small sella is usually constricted in the middle and the lancet is usually long and has a slender tip. (Fig. 30). – *Rhinolophus ferrumequinum*.

Additional characters: One or three mental grooves in the lower lip (Fig. 28) (very often the two lateral ones are reduced). Cf-frequency 79 - 84 kHz.

Distribution in Europe: Occurs in southern and central Europe, northwards to southern England and Wales, the Netherlands and Poland. Present on most Mediterranean Islands.

Photographs: Fig. 4, 20 - 21 and 28 – 31.

▶ smallest of the horseshoe bats, FA < 43 mm (usually 36 – 41 mm), D5: 46 – 53 mm, D3: 51 – 57 mm, P4.1: 5.7 – 7.5 mm; P4.2: 12.0 – 14.2 mm. The tip of the sella (= lower saddle process) is distinctly longer than the connecting process and in profile tapering to a point (Fig. 33 - 34). The fur is soft and sparse, grey on the back in younger individuals and brownish in older ones. – *Rhinolophus hipposideros*.

Additional characters: The lower lip has one mental groove (Fig. 32). Cf-frequency 108 - 115 kHz.

Distribution in Europe: It has the northernmost limit of distribution of all Rhinolophidae in Europe, reaching Ireland, the Netherlands, Thuringia in Germany and Poland. Common in the Mediterranean area and present on most islands.

Taxonomical note: The taxonomic position of some forms within this group is still not solved satisfyingly, especially in Asia. There are some morphological differences between the nominate form distributed over most of Europe and the populations from Northern

Africa (*R. h. escalerae*), Sicily, Crete (*R. h. minimus*), Cyprus and Western Anatolia.

Photographs: Fig. 32 – 34.

3) Second phalanx of the fourth finger (P4.2) less than twice as long as the first (P4.1) (P4.1: 7.6 – 9.2 mm; P4.2: 14.3 – 17.4 mm) (Fig. 38). Tip of the sella (= lower saddle process) narrow when viewed from the front and lower part not rounded (Fig. 37). The sella is wedge shaped when viewed from below. The horizontal furrow below the lancet is usually clearly indented in the middle when viewed from the front (Fig. 37). Connecting process (= upper saddle process) relatively long and straight, never curving downwards (Fig. 35 - 36). Bases of the hair whitish (tousled parts of the pelage have therefore a very light appearance), the tips of the hair are brown or greyish, often with a yellowish tinge (Fig. 35). FA 43.9 – 50.1 mm, D5: 54 – 62 mm, D3: 63 – 74 mm. – *Rhinolophus blasii*.

Additional characters: Cf-frequency 93 – 96 kHz.

Distribution in Europe: Restricted to south-eastern Europe from the northern Adriatic coast eastwards to Romania and all over the Balkans and Greece. Found also on some Greek islands including Crete.

Photographs: 35 – 38.

► Second phalanx of fourth finger (P4.2) more than twice as long as the first (P4.1) (Fig. 42). Tip of the sella (= lower saddle process) broad when viewed from the front and its lower margin rounded (Fig. 41 and 45). – **4**

4) Lancet narrows more or less gradually to its tip, it has only a slight constriction above the middle and the tip is broadly rounded (Fig. 41). Connecting process (= upper saddle process) is slightly horn-shaped, being pointed in profile and forward curving (slightly downwards) (Fig. 40). FA usually < 50 mm (44.0 – 51.0 mm), D5: 52 – 63 mm, D3: 63 – 76 mm, P4.1: 5.7 – 8.2 mm; P4.2: 16.4 – 18.1 mm. Coloration of the belly more greyish (Fig. 39), not as whitish as in *R. mehelyi*. The boundary between the back and underside is indistinct. – *Rhinolophus euryale*.

Additional characters: The antitragus of the ear (horizontal lobe at the base of the ear) is about half as high as the conch, its width is about equal to its height and it is only weakly indented close to the connection to the ear. Cf-frequency 102 – 107 kHz. Body mass usually around 12 g (9 – 15 g) in summer.

Distribution in Europe: Widest distributed species of the three medium sized horseshoe bats in the whole Mediterranean area and the Balkans, extends north to central France, northern Italy, Slovakia and Romania. Present on Corsica, Sardinia and Sicily but absent from the Balearic Islands.

Photographs: 39 – 42.

► Lancet is abruptly narrowed above the middle to a distinctly linear tip (Fig. 43 and 45). Connecting process (= upper saddle process) relatively blunt in profile and only slightly longer than the lower process (Fig. 44). FA usually > 49 mm (48.2 – 54.8 mm), D5: 57 – 67 mm, D3: 71 – 83 mm, P4.1: 6.5 – 9.3 mm; P4.2: 17.4 – 21.5 mm. Whitish belly coloration and clear boundary between the back and underside coloration in adult individuals. – *Rhinolophus mehelyi*.

Additional characters: The antitragus of the ear (horizontal lobe at the base of the ear) is not half as high as the conch, its width is greater than the height and it is strongly indented close to the connection to the ear, forming a well visible dent. Cf-frequency 106 - 112 kHz. Body mass usually around 15 g (12 – 18 g) in summer.

Some individuals from Sardinia have a distinct reddish brown to orange red coloration of the fur, even the belly can be coloured red in such individuals.

Distribution in Europe: Distributed in the Mediterranean area from central and southern Iberia, southern France, Sardinia, Sicily to Greece, range extends in the Balkans northwards to Romania.

Photographs: 19 and 43 – 45.

Plate 6: The five European species of horseshoe bats (Rhinolophidae).

R. ferrumequinum (28 - 31), *R. hipposideros* (32 - 34), *R. blasii* (35 - 38),
R. euryale (39 - 42), *R. mehelyi* (43 - 45).



Plate 6: The five European species of horseshoe bats (Rhinolophidae).

B) MOLOSSIDAE

The only European species, *Tadarida teniotis*, is very large and has long and broad ears projecting forward over the face (Fig. 16). The ears touch in front at their base. The posterior margin of the ear is broadened and has a conspicuous antitragus (Fig. 17). The long muzzle has usually five creases in the upper lip (Fig. 16). The fur is short and silky and has the structure of a mole's pelage. The back is blackish-grey, some individuals have a brownish tinge. At least one third of the tail extends beyond the tail membrane and has stiff hairs at the end (used as tactile organs) (Fig. 18). The wings are long and narrow. FA: 57.2 – 64.1 mm, D5 > 55 mm, D3 > 100 mm. –

Tadarida teniotis.

Additional characters: Thumb and especially the hind foot with white and curved bristles which are used like a comb to clean the fur. The high and fast flying bats emit well audible and loud sounds (8 - 11 kHz).

Distribution in Europe: Distributed in the Mediterranean region northwards to the southern Alps, Adriatic coast and Bulgaria. Present on most of the Mediterranean Islands.

Photographs: 16 – 18.

C) MINIOPTERIDAE

The single European species, *Miniopterus schreibersii*, has a very short muzzle and a humped forehead (Fig. 25). The ears are short and triangular and do not extend over the top of the head (Fig. 26) which has a dense, short and erected pelage reaching the back of the nose. The dorsal pelage is greyish-brown, sometimes brown or blackish. The underside is of a slightly lighter grey hue. The wings are very long and narrow and at rest the third and fourth fingers are bent to the inside in the joint between 1st and 2nd phalanges. FA: 42.4 – 48.0 mm, D5: 48 – 56 mm, D3: 78 – 89 mm. – *Miniopterus schreibersii*

Additional characters: In south-eastern Europe most individuals have a contrasting cinnamon coloured throat patch (Fig. 25) and forehead during moulting. Penis thin and long.

Distribution in Europe: The species occurs in southern and south-eastern Europe and is widely distributed in the Mediterranean and the Balkans and reaches as far north as Switzerland, southern Germany (currently extinct), Alsace in France and Slovakia.

Taxonomical note: Until recently, the Miniopteridae were treated as a subfamily of the Vespertilionidae but genetic studies have shown a degree of distinctiveness supporting a family level of its own. Morphological and physiological characters (like suspension of embryonic development) give further support to this classification.

Photographs: 25 – 27.

D) VESPERTILIONIDAE

1) only one pair of upper incisors, ears short and rounded, inside densely covered with hairs. Tail membrane heavily furred. Yellowish-brown pelage with frosted white tips, throat buffy-yellow. FA: 42 - 59 mm. – *Lasiurus cinereus*

Additional characters: Two pairs of teats.

Distribution in Europe: Very rare vagrant from North America, only a few records on Iceland and the Orkneys.

Photographs: none

► ears not covered with dense hairs, two pairs of upper incisors (but the second incisor might be hidden sometimes in the gum in *Hypsugo savii* or can be very small in *Pipistrellus kuhlii*). – **2**

2) ears connected in front at their base by a fold of skin (Fig. 46) and touching each other when erected (Fig. 48). Nostrils open above (Fig. 53 l.). – **3** (Subfamily Plecotini)

► ears widely separated in front, no fold of skin between the ears (ears separated by normal pelage instead) (Fig. 47 and 49). Nostril open to the front (Fig. 47). – **4**

3) ears over 30 mm long with numerous horizontal furrows (Fig. 48 and 50), folded at rest (Fig. 46). Spur without post calcareal lobe. – Genus *Plecotus* – see separate key to the species.

► ears shorter (up to 18 mm long) and wide with 5 - 6 furrows (Fig. 51 and 52). Ears never folded at rest. Dorsal fur blackish with light tips (Fig. 53), appearing frosted. FA: 36.5 – 43.5 mm, D5: 47 – 54 mm, D3: 63 – 71 mm. – *Barbastella barbastellus*

Additional characters: Ears often with a button-like process in the middle of the outside margin (Fig. 51 and 52). Spur with a post calcareal lobe divided by a keel.

Distribution in Europe: Southern and Central Europe, northwards to Britain, Scandinavia and Latvia, missing or rare in the southernmost parts of Europe, being there confined to the mountains. Present on the Balearic Islands, absent from Crete.


Photographs: 51 – 53 (right).



46



47



48



49



50



51



52



53

P. kolombatovici (46, 48),
E. nilssonii (47),
M. bechsteinii (49),
P. sardus (50),
B. barbastellus
 (51, 52, 53 r.),
P. auritus (53 l.)

4) Tragus long, pointed spear-shaped (Fig. 54). No post calcareal lobe at the spur (Fig. 57) (in whiskered bats and some Bechstein's bats a narrow edging of skin is present, but the tragus is tapering and reaches half of the ear length). Three premolars in the upper and lower jaw. – 5 (Genus *Myotis*)

► Tragus short, curved, with a rounded tip (Fig. 55), sometimes even with mushroom-shaped broadening (Fig. 56). Well developed post calcareal lobe at the spur (Fig. 58 - 61). Only one or two premolars. – 10

5) Big bats, FA > 50 mm. – Large *Myotis*-species – see separate key to the species.

► Smaller bats, FA < 50 mm. – 6

6) very big ears (Fig. 62), more than 20 mm long (21 - 26 mm), when folded forward projecting by nearly half their length beyond the tip of the snout (extending more than 8 mm) (Fig. 64 and 65). Wing membrane inserted at the base of the first toe (Fig. 66). FA 39.0 – 47.1 mm, D5: 50 – 57 mm, D3: 61 – 69 mm. – *Myotis bechsteinii*

Additional characters: Outside margin of the ear with 9 – 11 horizontal creases (Fig. 62 and 64). Spur straight (Fig. 67), sometimes with a narrow edging of skin.

Distribution in Europe: Distributed all over Europe but rare in the north (missing in most parts of Britain), in Scandinavia only in southern Sweden. Local in the southernmost parts of Europe. Absent from the Balearic Islands, Sardinia and Crete.

Photographs: 49 and 62, 64, 65, 67 and 72.


► ears less than 20 mm long (usually less than 18 mm), when folded forward extending the snout at most by 5 mm (Fig. 66). Maximum of 8 horizontal creases at the outside of the ear. – 7

Plate 8: Characters of the genus *Myotis* in comparison to other genera.

M. myotis (54), *E. serotinus* (55), *N. leisleri* (56, 60), *M. daubentonii* (57),
H. savii (58, 59), *N. noctula* (61).

Plate 9: The “long-eared” *Myotis*-species *Myotis bechsteinii* and *Myotis nattereri*.

M. bechsteinii (62, 64, 65, 67, 72), *M. nattereri* (63, 66, 68, 69, 71).
M. daubentonii (70)





7) Spur very long (longer than half of the margin of the tail membrane) and curved into S-shape (Fig. 68). Free margin of the tail membrane covered thickly with short, curved bristles (Fig. 69). Ears long with a very long (longer than half of the ear) and spear-shaped (lanceolate) tragus, ears light in colour (Fig. 63). FA: 34.4 – 44.0 mm, D5: 48 – 58 mm, D3: 65 – 74 mm. – *Myotis nattereri*

Additional characters: Wing inserted at the base of the outer toe (Fig. 68). Ventral pelage white (Fig. 71).



Distribution in Europe: Distributed all over Europe, in the north to Scotland and southern Scandinavia. Perhaps missing on Sardinia and Crete.

Photographs: 63, 66, 68, 69 and 71.

► spur straight or only slightly curved to one side (never S-shaped) and not longer than half of the margin of the tail membrane (Fig. 70), but there might be a small lobe at two third of the length of the margin of the tail membrane imitating the end of the spur (Fig. 70). Ears shorter. – **8**

8) very big hind foot with long bristles (Fig. 75), the hind foot length (HF) is larger than half of the tibia length (Tib) (Fig. 73). The wing membrane is inserted at the middle or base of the sole of the hind foot (Fig. 75) or at the tibia. The spur is about a third of the length of the tail membrane, but at two-thirds to three-quarters of length of the membrane is a distinct break, looking like the tip of the spur (therefore sometimes called the terminal lobe) (Fig. 77). Posterior margin of the ear without a distinct indentation (Fig. 79). – *Trawling Myotis* – see separate key to the species.

► hind foot smaller (Fig. 76), the hind foot length (HF) is only half, or less than half of the tibia length (Tib) (Fig. 74). Wing membrane inserted at the base of the outer toe (Fig. 76). Spur length is no more than half the length of the margin of the tail membrane and there is no terminal lobe or break present (Fig. 78). Posterior margin of the ear with a distinct indentation (Fig. 80). – **9**



M. capaccinii (73),
M. aurasceus (74, 76, 78, 80),
M. daubentonii (75, 77, 79)

9) FA usually > 37 mm (36.1 – 44.7 mm), D5: 49 – 58 mm, D3: 59 – 71 mm. The distinct indentation (notch) on the outside of the ear is nearly a right-angle (Fig. 82). The indentation is not reached by the tip of the tragus (Fig. 81). Spur without post calcareal lobe or any edging of skin. Dorsal fur long, woolly and especially in adult individuals with a distinct reddish tinge (Fig. 88) (the bats of the population on Sardinia are much darker blackish-brown without any reddish tinge, Fig. 89). – *Myotis emarginatus*

Additional characters: The free margin of the tail membrane sometimes with evident fringe but usually with sparse, short soft hairs which sometimes are hardly visible and might even be absent. The skin of the testes and the epididymids are pigmented dark black, even in older (several years old) males (Fig. 86 and 87).

Distribution in Europe: All over southern and Central Europe, northernmost occurrence in the Netherlands and southern Poland.

Taxonomical note: The population on Sardinia might represent a distinct subspecies as it differs remarkably from mainland populations in fur colour (Fig. 87 and 89).

Photographs: 3 and 81 – 89.

► smaller species, FA mostly < 36 mm (only a few individuals reach nearly 38 mm). Tragus extends beyond the indentation on the posterior margin of the ear (Fig. 80) (if not reaching the upper margin of the indentation, individuals are very small with FA < 33 mm: *Myotis alcathoe*). Pelage long and frizzy. Hair with dark bases and lighter tips, frequently with golden gloss. Spur often with narrow edging of skin (post calcareal lobe). The skin at the testes is never dark pigmented and at the epididymids it is only dark in young males, later it is light coloured. – *Whiskered bats* – see separate key to the species.

Plate 11: Characters of *Myotis emarginatus*.

M. emarginatus (81 - 89), 87 and 89 show an adult male from Sardinia.



10) smaller species, FA < 38 mm. Tragus rounded, club shaped. Posterior margin of the ear connected with the corner of the mouth by a narrow furrow only. – **11**

► bigger species, FA > 38 mm. – **12**

11) Last 1 or 2 tail vertebrae extending beyond tail membrane by 4 – 5 mm (Fig. 95). Post calcareal lobe narrow and always without keel (Fig. 96), sometimes even absent. Ears, face and wing membranes dark black (Fig. 90 - 92), much darker than in any European *Pipistrellus* species. Dorsal pelage long and dark with contrasting light golden tips, dorsal pelage contrasting to white ventral pelage (Fig. 90) (in adult individuals; younger ones are more uniformly brownish or dark greyish without lighter tips of the pelage (Fig. 91)). Tragus short and slightly broadening above (Fig. 92). Length of the front margin of the tragus almost corresponding to its greatest width (Fig. 93). Tip of the ear broadly rounded. FA: 31.4 – 37.9 mm, D5: 38 – 47 mm, D3: 52 – 63 mm. – *Hypsugo savii*

Additional characters: Tragus sometimes with two superimposed serrations at the base of the outside margin (Fig. 92). Characteristic penis morphology differing from all pipistrelles: penis relatively small and distal part slightly broadened. Between proximal and distal part characteristic right-angled bend (Fig. 94). Upper side of the penis with a medial groove. Upper canine (C¹) and second upper premolar (P⁴) in contact (Fig. 178).


Distribution in Europe: Mainly distributed in the south, northwards to Switzerland, Slovakia and the Danube. Vagrants further northwards. Present on most Mediterranean islands.

Photographs: 22, 58 – 59, 90 – 96 and 152 - 155. Drawings: 178.

► Last tail vertebrae extending to a maximum of 1 - 2 mm beyond the tail membrane (Fig. 99). Post calcareal lobe broad with well developed keel (Fig. 98). Tip of the ears narrowly rounded (Fig. 97). Dorsal and ventral pelage not contrasting. Ears and face usually not dark black (Fig. 97). – Genus *Pipistrellus* – see separate key to the species.

Plate 12: Characters of the genus *Hypsugo* and genus *Pipistrellus*.

H. savii (90 - 96), *P. nathusii* (97), *P. pipistrellus* (98, 99).



12) posterior margin of the ear with a narrow furrow extending towards the corner of the mouth but ending before it (Fig. 100). Tragus clearly longer than wide (Fig. 101). Post calcareal lobe narrow and usually without visible keel (rarely visible in *E. nilssonii*) – Genus *Eptesicus* – see separate key to the species.

▶ posterior margin of the ear with a broad furrow extending down below the line of the corner of the mouth and ending by it (Fig. 103, 104 and 107). Post calcareal lobe broad with well visible keel. – **13**

13) Tragus broadens above into mushroom shape (Fig. 104). Underside of the wing membrane adjacent to the forearm covered with short brown hair, underside of the wing close to the body covered with fur as well (as far distal as the line connecting the knee to the elbow, in *N. lasiopterus* even further). Ventral pelage only slightly lighter than dorsal one. Wing membrane inserted at the heel. D5 only a little bit longer than the 3rd and 4th metacarpals. Females with two teats. – Genus *Nyctalus* – see separate key to the species.

▶ The tragus is short, widening above, but not broadened above into mushroom shape (Fig. 107). Fine grey hair is only present on the underside of the wing along the forearm (only visible when the wing is closed). Dorsal pelage dark brown or blackish at the base and appearing frosted because of white hair tips (Fig. 106). Underside whitish or white and sharply demarcated from the dorsal side, in younger individuals and some adults more grey. Wing membrane inserted at the base of the first toe. FA: 40.8 – 50.3 mm, D5: 48 – 53 mm; D3: 69 - 76 mm. – *Vespertilio murinus*

Additional characters: Females with four teats. Penis long and very narrow (Fig. 108).

Distribution in Europe: Distributed in Central and eastern Europe northwards to central Scandinavia, westwards to Belgium and eastern France, southwards to Bulgaria and Greece. Absent from western France, Iberia, most of Italy, Peloponnese and all the Mediterranean islands. Photographs: 106 – 108.

Plate 13: Characters of the genera *Eptesicus*, *Nyctalus* and *Vespertilio*.

E. serotinus (100 - 102), *N. leisleri* (103 - 105), *V. murinus* (106 -108).



SEPARATE KEYS TO SPECIES GROUPS**I) Large *Myotis* species (*Myotis myotis* – *Myotis blythii* – *Myotis punicus*)**

The two widely distributed species *Myotis myotis* and *Myotis blythii* usually can be distinguished quite well. The third species, *Myotis punicus*, is much harder to identify, but it seems to occur in Europe only in strict allopatry on the islands of Corsica, Sardinia, Malta and Gozo. It shows a mixture of characters found in the other two species and its systematic position has long been unclear. As it is clearly separated from the other two species by its genetics (cytochrome b and microsatellites) its species status may be accepted.

▶▶▶ big bat, FA 55.0 – 66.9 mm, D5: 67 – 84 mm, D3: 89 – 107 mm. Muzzle heavily built (Fig. 109). Ears broad: > 16 mm and long: > 24.5 mm (24.4 – 27.8 mm) (Fig. 112). The anterior margin of the ear clearly curves backward and the posterior margin usually has 7 – 8 horizontal creases (Fig. 112). The tragus is broad at its base and usually has a small dark spot at its tip (Fig. 115). – *Myotis myotis*

Additional characters: CM³ > 9.4 mm (9.2 – 10.6 mm).

Distribution in Europe: Throughout Europe to the North Sea and Baltic Sea, being extinct on the British Isles. Vagrants to southern Sweden and Latvia. On the Azores, the Balearic Islands and Sicily. Contrary to older publications missing on Corsica, Sardinia and Malta.

Photographs: 54, 109, 112 and 115.

▶▶ slightly smaller bat, FA: 50.5 – 62.1 mm, D5: 63 – 81 mm, D3: 85 – 103 mm. Ears narrow: earW < 16 mm and shorter: earL < 24.5 mm (21.0 – 24.3 mm) (Fig. 113). The front margin of the ear curves backward less strongly and the ear tapers to more of a point, the outside margin usually has 5 – 6 horizontal creases (Fig. 113). Tragus is narrow at its base, spear-shaped (lanceolate) (Fig. 116) and reaches half of the ear length. Usually underside brighter white than in *M. myotis*. – *Myotis blythii*

Additional characters: Having a much more graceful appearance than the heavily built *M. myotis* and a more 'open' appearance of the face due to shorter muzzle and clearer skin (Fig. 110). Individuals from Switzerland usually have a whitish tuft of hairs between the ears. This white spot is present quite irregular in most other populations (for example in south-eastern Bulgaria and Turkey only 5 - 45 % of the individuals have the white tuft.). *M. myotis* never has this white spot. $CM^3 < 9.4$ mm (8.1 – 9.4 mm).

Distribution in Europe: Common in the Mediterranean part of Europe from Portugal to Turkey, in the north to northern Switzerland, Slovakia and the Czech Republic. On Sicily and Crete and many Greek islands. Contrary to older publications missing on Corsica, Sardinia and Malta.

Taxonomical note: The subspecies *M. b. omari* is paler and slightly bigger than the European subspecies *M. b. oxygnathus*: FA: 54.0 - 62.4 mm, CM^3 : 8.6 – 9.5 mm and occurs on Crete, other Greek islands and Cyprus. Recent genetic analyses placed *omari* closer to *M. myotis* than to *M. blythii*, but the systematic resolution within the large *Myotis* is quite low up to now. Further research is necessary.

Photographs: 54, 110, 113 and 116.

► Big bat, in size nearly as large as *M. myotis*: FA: 56.0 – 62.4 mm, D5: 73 – 74 mm, D3: 92 – 94 mm. The ears appear to be the largest and widest in relation to head size of all large *Myotis* bats (Fig. 111): EarL: 26.1 - 29.0 mm, earW: 14.7 - 17.9 mm. The ears are very wide in the medium part and as a consequence oval shaped (Fig. 114). Inside the long ears are 7 - 10 horizontal creases (Fig. 114). The tragus shape is quite variable, sometimes wide at its base and comparable to the one of *M. myotis* but lacks a black spot at the tip and its inward curve is slightly more developed, sometimes more like that of *M. blythii* (Fig. 117). – *Myotis punicus*

Additional characters: The muzzle and size of the eye are comparable to *M. blythii* (Fig. 111). Like in *M. blythii* clear line of demarcation between dorsal and ventral pelage coloration, ventral pelage white. Length of the upper tooth row between the two other species: CM^3 : 8.9 – 10.0 mm.

Distribution in Europe: In Europe only on Sardinia, Corsica, Malta and Gozo, there like in Northern Africa, the only large *Myotis* species. No zone of overlap in distribution with *M. myotis* or *M. blythii* known.

Taxonomical note: The populations on the Mediterranean islands might belong to a subspecies different from the African form.

Photographs: 12, 111, 114, 117.

Plate 14: Characters of the large *Myotis*: *M. myotis*, *M. blythii* and *M. punicus*.

M. myotis (109, 112, 115), *M. blythii* (110, 113, 116),

M. punicus (111, 114, 117).




II) Whiskered bats (*Myotis alcathoe* - *Myotis mystacinus* – *Myotis aurascens* & *Myotis brandtii*)

The identification of living whiskered bats, especially of young individuals and females, is difficult. A correct identification is sometimes possible only by using characters of the skull and teeth - if at all. It is still unclear, if the relatively big and lighter coloured whiskered bats of south-eastern Europe really deserve species rank (*Myotis aurascens*) or if they are just a subspecies of *M. mystacinus*. In addition it is not clear if they really belong to the taxon *aurascens*, initially described from the Caucasus.

With this key it is not possible to assign all individuals unambiguously to one species described at present, as the variability within *M. alcathoe* is little known and *M. mystacinus* and *M. aurascens* are very similar. Characters of *M. mystacinus* as given in this key refer to Central European populations. The subspecies *M. m. occidentalis* from Iberia is bigger and lighter in dorsal coloration. It is not possible to distinguish with this key alone between *M. aurascens*, *M. m. occidentalis* and *M. mystacinus* from regions other than Central Europe. We recommend therefore to treat *M. mystacinus* and *M. aurascens* in any species lists as one group and to specify it with its geographical origin. Determination is possible with molecular methods in *M. brandtii* and *M. alcathoe*, but the markers tested up to now do not resolve within *Myotis mystacinus* / *aurascens* / *occidentalis*.

Plate 15: Characters of Brandt's bat and whiskered bats (genus *Myotis*).

M. brandtii (118 - 121), *M. alcathoe* (122 – 125), *M. mystacinus* (126 - 129),
M. aurascens (130 - 133).

Plate 15: Characters of Brandt's bat and whiskered bats (genus *Myotis*).

1) small whiskered bat, FA usually < 32.8 mm, (30.8 – 33.3 mm, rarely > 33.0 mm), D5: 37 – 44 mm, D3: 50 – 56 mm. Ears short, the upper margin of the distinct indentation (notch) on the posterior margin of the ear is not reached by the tragus (Fig. 123) or hardly exceeded. Short thumb, D1 < 4.7 mm (3.8 – 4.7 mm, usually < 4.5 mm), Tib < 14.8 mm (13.5 – 14.8 mm, usually < 14.5 mm) and HF < 5.6 mm (5.1 – 5.6 mm). – *Myotis alcathoe*

Additional characters: Penis small with no marked thickening at the end (Fig. 124 and 125). The species resembles on the first view in general appearance, coloration and proportions *M. daubentonii* or *M. brandtii*, but is much smaller (Fig. 122). Hair of the back is reddish, only rarely with golden gloss. Young bats up to an age of one year and some older bats are more greyish. It has the shortest muzzle of all whiskered bats and the face of adults is pink to rufous, like *M. daubentonii*. The nostrils have a variable shape but the lateral part of the nostril is very well developed, the nostrils have therefore the shape of a heart (Fig. 122). The skin around the calcar is usually very light.

Distribution in Europe: Only sketchy information is available at present, records from Greece, Bulgaria, Hungary, Slovakia, Switzerland, France and Spain.

Photographs: 122 – 125.

► medium sized to big whiskered bat (usually FA > 33 mm), ears relatively long with a distinct indentation (notch) on the posterior margin of the ear. The long tragus projects above the indentation (Fig. 119, 127 and 131). Relatively long thumb (D1 > 4.3 mm, usually > 4.7 mm), Tib > 14.6 mm (usually > 15.3 mm) and HF > 5.8 mm (usually > 6.0 mm). – **2**

2) penis distinctly thickened at the end (club shaped) even in subadult males, but most obvious in adult ones (Fig. 120 and 121). Upper second premolar (P^3) is located within the tooth row and is rather large (about 2/3 of the size of the first upper premolar (P^2)) (Fig. 136), cingular cusp of the third upper premolar (P^4) is higher than the second premolar (P^3) (Fig. 136). Paraconuli usually present. FA: 33.0 – 38.2 mm, D5: 40 – 49 mm, D3: 48 – 61 mm. – *Myotis brandtii*

Additional characters: Nostrils usually heart-shaped (Fig. 118). The dorsal pelage in adults with a golden gloss. All bare parts of skin are medium to light brown (Fig. 118), not blackish-brown. The base of the ear and the tragus are clearly lighter (Fig. 119) (all colour characters of adults are quite similar in *M. aurascens*; in young individuals coloration resembles *M. mystacinus*!). A narrow post calcareal lobe is usually present.

Distribution in Europe: Distributed mostly in central and northern Europe, getting more rare to the south. It is absent from Ireland, Iberia, western France, Greece and all Mediterranean islands. In the Balkan countries confined to the mountains.

Photographs: 118 – 121.




Plate 16: Teeth characters of Brandt's bat and whiskered bats.

M. aurascens (134, 137), *M. mystacinus* (135, 138) *M. brandtii* (136, 139).

► the penis is thin, of equal width, not or only slightly thickened at its end (Fig. 128 – 129 and 132 – 133). Upper second premolar rather small (maximum 1/2 of the size of the first upper premolar) (Fig. 135) and sometimes displaced palatally of the tooth-row (Fig. 134). Cingular cusp of third upper premolar is small or absent, always lower than the second upper premolar (Fig. 134 and 135). Paraconuli usually absent. – 3 (see note above!).

3) slightly smaller, penis slightly thinner and shorter (Fig. 128 and 129). Ears usually dark blackish-brown without lighter colour inside (Fig. 127). The muzzle is dark black-brown (Fig. 126). Nostril not heart-shaped, lateral part often very narrow (Fig. 126). Dorsal fur coloration brown but without golden tips or golden gloss. Adult individuals in most cases with pronounced yellowish-brown hair at and around the sides of the neck, forming a well visible sort of ruff (Fig. 126). Dimensions of thumb (D1: 4.3 – 5.9 mm, usually < 5.3 mm), lower leg (Tib: 14.6 – 16.8 mm, usually < 16.2 mm) and hind foot smaller (HF: 5.8 – 7.4 mm, usually < 6.8 mm). FA: 32.0 – 36.5 mm, D5: 38 – 46 mm, D3: 48 – 58 mm. – *Myotis mystacinus*

Additional characters: The margin of the wing between the fifth finger and the leg of the same colour as the wing, not lighter. Upper second premolar small but usually within the tooth-row and about 1/3 to 1/2 of the size of the first upper premolar (Fig. 135).

Distribution in Europe: Distributed from central Scandinavia to Iberia. Range in Italy and the Balkans largely unknown due to possible confusion with *M. alcaethoe* and *M. aurascens*.

Taxonomical note: Characters given here refer to the central European populations; the subspecies *M. m. occidentalis* (Iberia) resembles in some of its external characters *M. aurascens*, but differs clearly in skull and dental morphology.

Photographs: 126 – 129. Drawings: 135 and 138.

► slightly bigger, penis relatively broader (Fig. 132 and 133). Ears brown, the inside of the ear and the base of the tragus lighter brown, sometimes even pinkish (Fig. 131). Nostril often heart-shaped, lateral part usually well developed (Fig. 130). In older individuals fur of the back with light golden tips. Adult individuals always without yellowish-brown hair on the sides of the neck, therefore ventral and dorsal colours of the fur sharply divided (Fig. 130). Large dimensions of thumb (D1: 5.2 – 7.0 mm, usually > 5.4 mm), lower leg (Tib: 15.7 – 18.1 mm, usually > 16.1 mm) and hind foot (HF: 6.8 – 8.7 mm, usually > 7.0 mm). FA: 32.0 – 37.4 mm, D5: 43 – 50 mm, D3: 52 – 61 mm. – *Myotis aurascens*

Additional characters: The margin of the wing membrane between the fifth finger and the leg usually has a very thin white or at least light border. Second upper and lower premolar very small, the upper one often dislocated palatally (1/4 to 1/3 of the size of the first upper premolar) (Fig. 134 and 137).

Distribution in Europe: Due to problems in species determination, the range is not well known in Europe. Common in Greece and Bulgaria, northwards to Romania and Serbia. Along the Adriatic coast to Northern Italy. Perhaps most of the populations in Italy and at least in parts of Hungary belong to this form as well.

Taxonomical note: see note at the beginning of the whiskered bat key. Up to now it has not been possible to distinguish *M. aurascens* and *M. mystacinus* by their genetics. So *M. aurascens* might only be a subspecies of *M. mystacinus* and is perhaps different from the true *aurascens* from the Caucasus. Another possible name for these bats is *Myotis mystacinus bulgaricus*.

Photographs: 130 – 133 (and 1 – 2). Drawings: 134 and 137.

III) Trawling *Myotis* (*Myotis daubentonii* – *Myotis capaccinii* – *Myotis dasycneme*)

1) fairly large species, FA > 42 mm (usually 43.0 – 49.0 mm), D5: 51 – 61 mm, D3: 72 – 77 mm. Tragus relatively short (shorter than half of the ear length) and for a *Myotis* species unusually short and broadly rounded at its tip (Fig. 149). Wing membrane inserted at the ankle of the foot (Fig. 151). Tail membrane with very fine whitish hairs on underside along lower leg up to the spur (Fig. 151). The pelage is dense and greyish-brown on the back and greyish-white on the underside (Fig. 148).

– *Myotis dasycneme*

Additional characters: Penis is widest at its base and tapers towards the tip (Fig. 150).

Distribution in Europe: It is found from north-eastern France along the coast of the North Sea through northern Germany to southern Scandinavia, along the Baltic Sea to the Russian Plain and in the south to Slovakia, Croatia, Hungary and Romania.

Photographs: 148 – 151.

► smaller, FA usually < 42 mm. - **2**

2) wing membrane inserted before ankle on the lower leg (tibia) (Fig. 147). Hind foot very big. Tragus long, reaches at least half of the ear length and is curved into slight S-shape (Fig. 145). Dorsal pelage a striking grey (Fig. 144). Tibia and tail membrane covered on dorsal and ventral side with downy hairs reaching from the leg to about the middle of the tail membrane (Fig. 147). The pelage on the back is a light smoky grey, rarely with a brownish tinge. The underside is grey. FA: 38.4 – 44.0 mm (but rarely more than 43.0 mm), D5: 48.6 – 56.4 mm, D3: 64 – 71 mm. – *Myotis capaccinii*

Additional characters: The nostrils are somewhat protruding giving the species a characteristic profile (Fig. 144). Penis slightly broadened towards its tip (Fig. 146).

Distribution in Europe: Distributed in the Mediterranean area and the Balkans. From the west coast of Spain to southern France, Italy, southern Switzerland, all the Balkan countries to Romania in the north. Present on all large Mediterranean islands.

Photographs: 144 – 147.

► wing membrane inserted between the ankle and the middle of the sole of the hind foot (Fig. 143), sometimes closer to the base of the first toe. Tragus long, reaches half the ear length, straight or slightly curved but not into S-shape (Fig. 141). Dorsal fur brownish, shiny. Tibia and tail membrane not hairy (Fig. 143). FA: 33.1 – 42.0 mm, D5: 39 – 52 mm, D3: 53 – 65 mm. – *Myotis daubentonii*

Additional characters: Penis parallel sided or slightly broadened to its tip (Fig. 142).


Distribution in Europe: It occurs throughout almost all of Europe, being absent only from northern Scandinavia, northern Scotland, Sicily, southern Greece and Crete. (In contradiction to earlier references largely overlaps in its distribution in the south with *M. capaccinii*.).

Taxonomical note: In Spain smaller and different coloured bats occur in sympatry with the typical form and were described as species of its own: *M. nathalinae*. But as individuals resembling the *nathalinae*-type were found in several parts of Europe as well and no genetic differences between those and typical *M. daubentonii* were found, *nathalinae* is currently regarded as a synonym of *M. daubentonii*.

Photographs: 57, 70 and 140-143.

Plate 17: Characters of the trawling *Myotis* (genus *Myotis*, subgenus *Leuconoe*).

M. daubentonii (140 - 143), *M. capaccinii* (144 - 147),
M. dasycneme (148 - 151).



IV) Genus *Pipistrellus* (*Pipistrellus pipistrellus* – *Pipistrellus pygmaeus* – *Pipistrellus kuhlii* – *Pipistrellus nathusii*)

Pipistrelles can be quite difficult to identify. It is important to consider the teeth and a set of other characters. However, with some experience most of the “qualitative” characters can be evaluated on first sight.

1) First upper incisor with a single cusp, second upper incisor small (without magnifying lens it appears to be only one single cusped tooth) (Fig. 179 and 182). The last upper premolar (P^4) is in contact with the canine (Fig. 179) (no small premolar (P^2) visible from the outside, if present, displaced to the inside). Usually a well defined white stripe along the margin of the arm wing membrane between fifth finger and hind foot (Fig. 183), usually also present along tail membrane and between fifth and fourth finger. Ears lighter brown, usually not blackish (Fig. 160 and 162). Dorsal pelage with dark black bases and sandy yellowish-brown tips (Fig. 160). Some individuals are dark brown without light tips, they resemble in coloration *P. nathusii*. FA: 30.3 – 37.1 mm, D5: 40 – 45 mm, D3: 54 – 61 mm. – *Pipistrellus kuhlii*

Additional characters: Posterior margin of the ear with a sharp indentation (Fig. 162).

Penis is spear-shaped and without a medial stripe (Fig. 163) (its colour and shape even in juveniles quite similar to *Nyctalus leisleri*). The white stripe along the wing membrane is about 1 - 2 mm wide but up to 5 mm wide in animals from the south of the range.

Distribution in Europe: Occurring mainly in southern Europe in the whole Mediterranean, extending range northwards, reaching recently the southernmost parts of Germany and Austria. Missing or rare in the northern Balkan countries, absent from Romania, confined to the south in Bulgaria.

Photographs: 160 – 163, 175 and 182 – 183. Drawings: 179 (and 174).

► First upper incisor with two cusps and second incisor also clearly visible (Fig. 180 and 181). Wing membrane without a well defined white margin but sometimes with slightly lighter whitish margin. – 2

2) bigger species, FA: 32.2 – 37.1 mm. D5 usually > 43 mm (41 – 48 mm), D3: 57 – 62 mm. Pelage extends considerably on the tail membrane up to its proximal half. First upper premolar (P^2) clearly visible from outside and within the tooth row (Fig. 181). Second upper incisor (I^3) longer than the lower cusp of the first upper incisor (I^2) (Fig. 181). Cell of the wing membrane between the first joint of the fifth finger and the elbow divided by a keel (sometimes absent in small individuals) (Fig. 172 and 173). Coloration of the dorsal pelage dark brown (Fig. 156). – *Pipistrellus nathusii*

Additional characters: Margin of the wing membrane usually with a diffuse yellowish-white margin between leg and fifth finger. Underside of the tail membrane hairy along the lower leg (tibia). Characteristic gap between the second and third lower incisors (between I_2 and I_3). Penis well differentiated from all the other European pipistrelles in being robust and ovoid with a medial groove and strong hairs (Fig. 159). (Contrasting to earlier references, the relation between thumbL and width of the wrist is not species specific.)

Distribution in Europe: Occurs in Eastern, Central and Southern Europe. Common along the Baltic Sea, southern Scandinavia and northern Germany. Also present in the Balkans and on Corsica. Migrating to Sardinia and Sicily and Iberia. In the southern range in summer predominantly males, during migration and winter time also females.

Photographs: 97, 156 – 159 and 173. Drawings: 172 and 181.

► smaller species, FA < 34.6 mm. D5 < 43 mm (usually < 41 mm). Cusp of the first upper premolar (P^2) visible between canine and second upper premolar (P^4) but displaced inside of the tooth row (Fig. 180) (sometimes barely or not visible from outside, but unlike *P. kuhlii* P^4 and C^1 not directly in contact). Second upper incisor (I^3) shorter than the lower cusp of the first upper incisor (I^2) (Fig. 180). The three cusps of the incisor are stepped and look therefore even without magnification like a three-toothed comb (Fig. 180). No hair on the underside of the tail membrane along

the tibia. Cell of the wing membrane between the first joint of the fifth finger and the elbow not divided by a keel (Fig. 174 - 177) (only some very big females of *P. pipistrellus* show sometimes a weak keel). – 3

3) slightly bigger species, FA: 29.2 – 33.5 mm (in extremes 28.0 – 34.5 mm), D5: 37 – 41 mm, D3: 50 – 56 mm. Without internarial ridge between the nostrils (Fig. 165, 184) (only very dehydrated individuals do sometimes show a weakly developed ridge). The muzzle is longer and gradually narrowing. The glandular bumps are white (Fig. 186) or sometimes whitish-grey. The length of the inner margin of the ear is longer (8 - 9 mm) (Fig. 166). The pelage is dense but not as smooth as in *P. pygmaeus* and dark brown on the back, often rusty, sometimes a paler medium brown. Ears and muzzle are mostly black (Fig. 164 and 166), but in the south of the range some adults have a pale area around the eyes. Face and ears differ strongly from the pelage by their dark coloration. The penis is dark grey to greyish-brown and the glans penis with a contrasting pale median stripe (Fig. 167, 189). The wing cell connecting the first joint of the fifth finger with the elbow is not divided by an additional keel (Fig. 174, 175 and 192), but the cell above this is quite short, it usually does not extend to the forearm (Fig. 174 and 175). – *Pipistrellus pipistrellus*

Additional characters: In most individuals the second phalanx of the 3rd finger (P3.2) is 1 - 3 mm longer than the 3rd phalanx (P3.3) (Fig. 194), but sometimes they are of similar length (P3.2: 7.9 – 8.9 mm, P3.3: 6.0 – 8.4 mm). There is usually no gap between the second and third lower incisors, they are in contact (I_2 and I_3). Terminal frequency around 45 kHz (43 – 49 kHz, in extremes 41 – 52 kHz).

Distribution in Europe: The species is distributed all over Europe to southern Scandinavia and the Baltic states. Its northern limit of distribution is further south than in *P. pygmaeus*. In some Mediterranean areas *P. pipistrellus* is rarer than *P. pygmaeus*, but in most central European countries *P. pipistrellus* is the most common and widespread bat.

Taxonomical note: Within this species or species group some taxonomic questions still require clarification: some populations in eastern Thrace, the Peloponnese and on some

Greek islands are quite delicately built and show intermediate characters to *P. pygmaeus* and emit echolocation calls with a terminal frequency around 50 kHz; but they never have a yellowish or orange penis and vagina. In contrast some individuals from Sardinia are of a more robust build and resemble much more *P. kuhlii*, but they always show the typical characters of the upper incisors.

Photographs: 98 – 99, 164 – 167, 184, 186, 188 – 189, 192 and 194.

Drawings: 174 and 180.

► slightly smaller, FA: 27.7 – 32.3 mm, D5: 33 – 40 mm, D3: 46 – 55 mm. Obvious internarial ridge between the nostrils (Fig. 169, 185). In dorsal view, the short muzzle is parallel sided for approximately two-thirds of its length, then converging. Glandular bumps during reproductive season obvious orange or yellow, throughout the year at least with an orange or yellow tinge (Fig. 187). The ears are shorter, the length of the inner margin is 7 – 8 mm (Fig. 170). Pale bald areas on the face, especially between the ears and the eyes and around the eyes (Fig. 168). Facial skin and ears are not darker in colour than the pelage coloration. Very dense silky pelage, dorsal reddish-brown, in winter more olive brown, underside yellowish-grey. In the south the summer pelage often sandy coloured. Ears lighter than in *P. pipistrellus*. Like in *P. pipistrellus* there is only one cell in the wing membrane between the first joint of the fifth finger and the elbow. In addition the next cell above (closer to the wrist) is also not divided in most individuals and connecting forearm and fifth finger (Fig. 176, 177 and 193). Penis in full adults with obvious yellow coloration, during reproduction time often orange (Fig. 190), glans penis always without paler medial stripe (Fig. 171). In subadults or juveniles the penis is whitish, often with a yellow tinge, never brownish and always without pale stripe. In females the skin around the vagina is also orange coloured (Fig. 191), at least when they are in oestrus. – *Pipistrellus pygmaeus*

Additional characters: The uropatagium is densely covered with hairs on its proximal third.

The second (P3.2) and the third phalanges (P3.3) of the third finger are more or less of

the same length (P3.2: 6.6 – 8.7 mm, P3.3: 6.3 – 8.2 mm) (Fig. 195). Obvious musk like odour, especially during mating season. Like in *P. nathusii* there is usually a gap between the second and third lower incisors (I_2 and I_3). Terminal frequency around 55 kHz (52 – 57 kHz, in extremes 50 – 64 kHz).

Distribution in Europe: The species seems to range all over Europe from Scotland and southern Scandinavia to Iberia and European Turkey, but records are missing from some regions like the northern Balkans and southernmost Italy. *P. pygmaeus* is more common in northern and southern parts of Europe, in Central Europe mostly confined to the valleys of larger river systems. The distribution is patchier than in *P. pipistrellus*.

Photographs: 168 – 171, 177, 185, 187, 190 – 191, 193 and 195.

Drawings: 176 (and 180).

Plate 18: Characters of the species of the genera *Hypsugo* and *Pipistrellus*.

H. savii (152 - 155), *P. nathusii* (156 - 159), *P. kuhlii* (160 - 163),
P. pipistrellus (164 - 167), *P. pygmaeus* (168 - 171).




Plate 19: Characters of the species of the genera *Hypsugo* and *Pipistrellus*.


P. nathusii (172, 173, 181), *P. pipistrellus* (174, 173, 180),
P. pygmaeus (176, 177), *H. savii* (178), *P. kuhlii* (179, 182, 183).

Plate 20: Characters of the species *Pipistrellus pipistrellus* and *P. pygmaeus*.

P. pipistrellus (184, 186, 188, 189, 192, 194),
P. pygmaeus (185, 187, 190, 191, 193, 195).







V) Genus *Eptesicus* (*Eptesicus serotinus* – *Eptesicus nilssonii* – *Eptesicus bottae*)

The two widely distributed species *E. serotinus* and *E. nilssonii* can be separated by external measurements only. In Europe *E. bottae* occurs only on some of the Greek islands off the Anatolian coast.

▶▶▶ large and robust species, FA: 48.0 – 58.0 mm, D5 > 60 mm (59 – 69 mm all over Europe, 61 – 67 mm in southern Greece and the Greek islands), D3: 84 – 92 mm. The muzzle is wide and robust (Fig. 199). The tragus is broad and the ears are long (Fig. 200). The penis is only slightly widened towards the end and its upper side has a weak medial ridge (Fig. 201). Pelage on the back ranging in colour from dark brown to sometimes yellowish or golden brown (Fig. 199). Dark black ears and muzzle. – *Eptesicus serotinus*

Additional characters: In south-eastern Europe, especially on the Greek islands, the dorsal pelage is a light yellowish brown. Hair on the back long (about 11 mm).

Distribution in Europe: Ranging all over Europe, in the north to central Britain, the southernmost Sweden and the Baltic states. Absent from Ireland and perhaps also from Sardinia.

Taxonomical note: A small serotine has been described as a species on its own from Romania: *Eptesicus sodalis* and has been subsequently found in various parts of Europe. The form *sodalis* is believed now to be a synonym of *Eptesicus serotinus*, as these bats seem to be just unusual small individuals.

Photographs: 55 and 100 – 102 and 199 – 201.

▶▶ smaller species, FA: 37.0 – 44.0 mm, D5: 45 – 56 mm, D3: 62 – 68 mm. Dorsal pelage dark brown to black with light golden tips on the back and the forehead (Fig. 196). A well defined line of demarcation along the sides of the neck towards the light yellowish brown underparts. – *Eptesicus nilssonii*

Additional characters: Usually small tufts of golden or yellowish hair at the front margins of the ears (Fig. 196). Ears shorter than in the other *Eptesicus* species (Fig. 197).

In principle this is an unmistakable species, but misidentifications do occur with *Hypsugo savii* (that is smaller, has a different ear and tragus and a characteristic penis with bend) or *Vespertilio murinus* (in *V. murinus* the silver hair tips do not reach the forehead, the pelage on the chin is white or at least light, it has well developed post calcareal lobe, its penis is long and very narrow. The flight membranes of *V. murinus* are greyish brown, but gleaming black in *Eptesicus*).

Distribution in Europe: Boreal species, with the most northerly range of all species (even reproduces north of the Arctic circle). Distribution gets more scattered to the south, being there mostly confined to the mountains. In the south-west to the Swiss Alps, in the south to the Dinaric Alps and Carpathians. One single record from Bulgaria.

Photographs: 47 and 196 – 197.

► medium sized species, FA: 43.3 – 50.0 mm (in extremes 37.6 – 52.1 mm). D5 < 60 mm (54 – 58 mm). Shorter muzzle and in relation bigger eyes (Fig. 202). Ears shorter and tragus more narrow (Fig. 203). Penis widely broadened towards the end and with a small triangular furrow at the tip (Fig. 204). Coloration similar to the light forms of *E. serotinus* yellowish-brown on the back. Ventral fur lighter than in *E. serotinus*, more whitish (Fig. 202). Dark black ears and face. $CM^3 < 7.0$ mm (in *E. serotinus* $CM^3 > 7.2$ mm). – *Eptesicus bottae*

Additional characters: Pelage on the back shorter, hair about 8 – 9 mm long.

Distribution in Europe: In Europe only on the Greek Islands off the Anatolian coast: Rhodes and possibly Samos and some more islands.

Taxonomical note: The characters outlined above are only valid for the Anatolian subspecies *E. b. anatolicus* distributed also on the Aegean Islands. The form *anatolicus* might be a distinct species as well.


Photographs: 202 – 203. Drawings: 204.

Plate 21: Characters of the species of the genus *Eptesicus*.

E. nilssonii (196, 197, (198)), *E. serotinus* (199 - 201),
E. bottae (202 - 204).



(198)



VI) Genus *Nyctalus* (*Nyctalus noctula* – *Nyctalus leisleri* – *Nyctalus lasiopterus*)

All three European species are well separated by their size and it is possible to determine them by using only the forearm length (FA).

▶▶▶ large species, FA: 48.0 – 55.4 mm (in extremes 47 - 59 mm), D5: 47 – 58 mm, D3: 85 – 98 mm. Pelage uniformly reddish-brown in late autumn and winter with a greyer tinge. – *Nyctalus noctula*

Distribution in Europe: Ranging throughout Europe except for Ireland, Scotland, northern Scandinavia and the southernmost parts of Greece and Italy. Mostly absent from the Mediterranean Islands.

Photographs: 61 and 208 – 210.

▶▶ very large species, FA: 64 – 68 mm (extremes 61 – 70 mm), D5: 69 – 74 mm, D3: 108 – 116 mm. Pelage uniformly reddish-brown. Very broad ears (Fig. 212), heavy muzzle (Fig. 211). – *Nyctalus lasiopterus*

Additional characters: Especially males with long lion-like fur on the neck (Fig. 211).

Distribution in Europe: Scattered records all over southern Europe, single records also in central Europe. Most common in Spain and Greece.

Photographs: 211 - 213.

▶ medium sized species, FA: 39.2 – 45.6 mm (extremes 38.0 – 47.1 mm), D5: 43 – 51 mm, D3: 70 – 78 mm. Pelage uniformly brown to dark brown without reddish tinge. Dorsal pelage two-coloured with darker bases. Ears narrower (Fig. 206). – *Nyctalus leisleri*

Additional characters: basal inner part of the ear and the fold of skin connecting the ear with the mouth are very often lighter in colour than the rest of the ear (Fig. 206).

Distribution in Europe: Records from all over Europe, but mostly absent from Scandinavia and Estonia. Missing in southern Italy, Sicily and Crete.

Photographs: 56, 60, 103 – 105, 205 – 207.

Plate 22: Characters of the species of the genus *Nyctalus*.

N. leisleri (205 - 207), *N. noctula* (208 - 210), *N. lasiopterus* (211 - 213).



VII) Genus *Plecotus* (*Plecotus auritus* – *Plecotus austriacus* – *Plecotus macrobullaris* – *Plecotus kolombatovici* – *Plecotus sardus*)

1) thumb short ($D1 < 6.5$ mm) (Fig. 232 and 238). Thumb claw usually less than 2 mm long (Fig. 232). Hind foot short (HF without claws < 8 mm). Hairs on the toes short and plain (Fig. 233 and 239). Penis thickened and rounded towards the end (Fig. 234 and 240). Protuberance above the eye small (in diameter smaller than diameter of the eye) (Fig. 217 and 223). – **2**

►► thumb long ($D1 > 6.5$ mm) (Fig. 229 and 235). Thumb claw long and curved (usually longer than 2 mm) (Fig. 235). Hind foot large (HF > 8 mm) with long hairs sticking up (Fig. 230 and 236). Tip of penis not thickened but penis parallel sided (Fig. 237 and 243) or tapered to the tip (Fig. 231). Protuberance above the eye medium-sized to big (Fig. 214 and 220). – **3**

► thumb of medium length ($D1: 6.0 - 6.5$ mm) (Fig. 241). Thumb claw long and curved (2.0 – 3.1 mm) (Fig. 241). Tragus length > 18 mm (Fig. 228). Hind foot small to medium-sized (HF: 6.7 – 7.7 mm). Long sticking hairs only at the toes, not at the foot (Fig. 242). $D5$ usually > 55 mm (minimum 54 mm), $D3$ usually > 71 mm (minimum 66 mm). Dorsal fur brown to greyish-brown, underside lighter but not white, sharp boundary. Penis parallel sided, only tapered at the tip (Fig. 243). FA: 40.9 – 42.3 mm. – *Plecotus sardus*

Additional characters: Facial parts in older bats light. Chin with an obvious round gland.

Distribution in Europe: Endemic to the island of Sardinia.

Photographs: 50, 226 - 228 and 241 – 243.

2) Small species: FA in males < 38 mm, in females < 39 mm (36.1 – 39.3 mm, in extremes 41.0 mm). $D3 < 65$ mm (61 – 66 mm). $D5 < 52$ mm (46 – 51 mm). Tib < 18

mm (15.2 - 18.3 mm). TragL usually < 14 mm, TragW usually < 5.2 mm. Dorsal pelage brown-grey to brownish. – *Plecotus kolombatovici*

Additional characters: $CM^3 < 5.7$ mm (> 5.7 mm in *P. austriacus*).

Distribution in Europe: In Europe only along the Adriatic coast, many Adriatic islands and Greece.

Taxonomical notes: BENDA et al. (2004) described a new form of long-eared bats from Northern Africa ("*gaisleri*"). Due to a lack of a sympatric occurrence of the three closely related forms "*teneriffae*" (from the Canary islands), "*gaisleri*" (northern Africa) and "*kolombatovici*" (Adriatic coast, Greece and Turkey) they were claimed to be subspecies of one single species. As *teneriffae* was the species described first, they refer to them as *Plecotus teneriffae teneriffae*, *P. t. gaisleri* and *P. t. kolombatovici*. Nevertheless, all three forms might represent independent species, as two of them (*P. t. gaisleri* and *P. t. kolombatovici*) were found in sympatry on the island of Pantelleria. Further research is needed to solve this taxonomical problem. Characters given in the identification key above are only valid for the form "*kolombatovici*" in Europe. *P. t. gaisleri* differs in having a darker coloration of pelage and naked parts and a larger thumb and forearm (FA: 37.2 - 40.9 mm). *P. t. teneriffae* is quite large (FA: 40.1 – 46.0 mm) and has a dark ash-grey ventral pelage coloration.

Photographs: 46, 48, 223 - 225 and 238 – 240.

► Larger species: FA in males usually > 38 mm, in females > 39 mm (36.5 – 43.5 mm). D3 > 64 mm (64 – 71 mm). D5 > 51 mm (48 – 55 mm). Tib > 18 mm. TragL: 14.0-16.0 mm and TragW > 5.4 mm. Dorsal pelage grey, but in east-mediterranean populations (Greece, Bulgaria, European part of Turkey sometimes brownish-grey). –

Plecotus austriacus

Additional characters: $CM^3 > 5.7$ mm (< 5.7 mm in *P. kolombatovici*).

Distribution in Europe: Distributed all over central and southern Europe, in the north to southern Britain, missing in Denmark and Scandinavia. Everywhere in the south including most Mediterranean islands, but missing on the Adriatic islands.

Photographs: 217 – 219, 232 – 234.

3) long upright hairs on the whole hind foot and toes (Fig. 230). TragL < 15.5 mm. D5 < 55 mm (47 – 56 mm), D3 < 66 mm. Dorsal pelage light brown, brown or reddish-brown. Ventral fur lighter, usually yellowish-brown. Usually without a clear boundary between dorsal and ventral coloration. Facial parts of the skin, ears and tragus brownish coloured without grey or black (Fig. 214). Without smooth triangular pad on the lower lip (Fig. 215). Protuberance above the eye big (1.5 - 2.0 mm in diameter). Penis tapered continuously from the base towards the tip (Fig. 231) (triangular shaped). FA: 35.5 – 42.5 mm. – *Plecotus auritus*

Distribution in Europe: Distributed throughout almost all of Europe but getting rarer to the south and often being confined to mountainous areas, missing on most islands except Sardinia.

Taxonomical note: the subspecies *Plecotus auritus begognae* from southern Iberia is larger (FA: 38.3 - 43.5 mm, TL: 5.9 – 7.6 mm, HF: 6.7 – 9.6 mm, TragL: 12.6 – 17.1 mm, TragW: 4.4 – 5.8 mm).

Photographs: 53 (left), 214 – 216 and 229 – 231.

► the hind foot is sparsely haired with long sticking hairs only visible at the toes (Fig. 236). Thumb and thumb claw shorter (Fig. 235) and FA usually bigger (FA 39.6 – 45.0 mm.) than in *P. auritus*. TragL usually > 16 mm. D5 > 51 mm. D3 > 63 mm. Pelage long and silky, dorsal brownish-grey to grey, ventral fur conspicuously white: ventral hairs with white tips and greyish proximal parts. A hard triangular pad extends towards the chin at the lower lip (Fig. 221), this pad is darkly pigmented at least in younger individuals. Penis parallel sided, only tapered at the tip (Fig. 237). – *Plecotus macrobullaris*

Distribution in Europe: Distributed in the Alpine parts of the Pyrenees, Corsica, Alps, the Dinaric Alps, Pindus mountains and Crete. Perhaps also present in the Carpathians and Balkan mountains (as it is distributed eastwards to the Caucasus and the Taurus mountains).

Taxonomical note: Currently two recognised subspecies: *P. m. alpinus* ranging from the Alpine regions, northern Italy and Croatia westwards and *P. m. macrobullaris* from Greece eastwards to Turkey, Armenia and Ossetia. Details of distribution are not known yet.

Photographs: 220 – 222 and 235 – 237.

Plate 23: Characters of the species of the genus *Plecotus*.

P. auritus (214 - 216), *P. austriacus* (217 - 219), *P. macrobullaris* (220 - 222),
P. kolombatovici (223 - 225), *P. sardus* (226 - 228).

Plate 24: Characters of the species of the genus *Plecotus*.

P. auritus (229 - 231), *P. austriacus* (232 - 234), *P. macrobullaris* (235 - 237),
P. kolombatovici (238 - 240), *P. sardus* (241 - 243).

Plate 23: Characters of the species of the genus *Plecotus*.

Plate 24: Characters of the species of the genus *Plecotus*.

REFERENCES

- AGUIRRE-MENDI, P. T., J. L. GARCIA-MUDARRA, J. JUSTE, & C. IBANEZ (2004): Presence of *Myotis alcathoe* HELVERSEN & HELLER, 2001 (Chiroptera: Vespertilionidae) in the Iberian Peninsula. – *Acta Chiropterologica*, 6 (1): 49-57.
- ARLETTAZ, R., M. RUEDI & J. HAUSSEER (1991): Field morphological identification of *Myotis myotis* and *Myotis blythi*: a multivariate approach. – *Myotis* 29: 7-16.
- ARLETTAZ, R., M. RUEDI, C. IBANEZ, J. PALMEIRIM & J. HAUSSEER (1997): A new perspective on the zoogeography of the sibling mouse-eared bat species *Myotis myotis* and *Myotis blythii*: morphological, genetical and ecological evidence. – *Journal of Zoology London* 242: 45-62.
- BENDA, P. & K.A. TSYTSULINA (2000): Taxonomic revision of *Myotis mystacinus* group (Mammalia: Chiroptera) in the western Palearctic. – *Acta Societatis Zoologica Bohemica*, 64: 331-398.
- BENDA, P. (2004): *Myotis auraszens* – Steppen-Bartfledermaus. – In: Krapp, F. (Hrsg.): *Handbuch der Säugetiere Europas, Fledertiere II*: 1149-1158.
- BENDA, P., A. KIEFER, V. HANAK & M. VEITH (2004): Systematic status of African populations of long-eared bats, Genus *Plecotus* (Mammalia: Chiroptera). – *Folia Zoologica* 53, Monograph 1: 47 pp.
- CASTELLA, V., M. RUEDI, L. EXCOFFIER, C. IBANEZ, R. ARLETTAZ & J. HAUSSEER (2000): Is the Gibraltar Strait a barrier to gene flow for the bat *Myotis myotis*? – *Molecular Ecology* 9: 1761-1772.
- DE PAZ, O. (1994): Systematic position of *Plecotus* from the Iberian Peninsula. – *Mammalia* 58 (3): 423-432.
- DE PAZ, O. (1995): Geographic variation of the greater horseshoe bat (*Rhinolophus ferrumequinum*) in the west-half of the Palearctic region. – *Myotis* 32/33: 33-44.
- FELTEN, H., F. SPITZENBERGER & G. STORCH (1977): Zur Kleinsäugerfauna West-Anatoliens, Teil IIIa. – *Senckenbergiana biologica* 58 (1/2): 1-44.
- GAUCKLER, A. & M. KRAUS (1970): Kennzeichen und Verbreitung von *Myotis brandti*. – *Zeitschrift für Säugetierkunde*, 35: 113-124.
- HÄUSSLER, U., A. NAGEL, M. BRAUN & A. ARNOLD (1999): External characters discriminating sibling species of European pipistrelles, *Pipistrellus pipistrellus* and *P. pygmaeus*. – *Myotis* 37: 27-40.
- HÄUSSLER, U. (2003): Kurzer Bestimmungsschlüssel mitteleuropäischer Fledermäuse nach äußeren Kennzeichen und Zahnmerkmale. – In: BRAUN, M. & F. DIETERLEN (Hrsg.): *Die Säugetiere Baden-Württembergs, Band 1: Allgemeiner Teil, Fledermäuse (Chiroptera)*: 333-339; Verlag Eugen Ulmer.

- HÄUSSLER, U. & M. BRAUN (2003): Mückenfledermaus *Pipistrellus pygmaeus*. – In: BRAUN, M. & F. DIETERLEN (Hrsg.): Die Säugetiere Baden-Württembergs, Band 1: Allgemeiner Teil, Fledermäuse (Chiroptera): 544-568; Verlag Eugen Ulmer.
- HILL, J.E. & D.W. YALDEN (1990): The status of the hoary bat, *Lasiurus cinereus*, as a British species. – Journal of Zoology London 222: 694-697.
- HORACEK, I. & P. BENDA (2004): *Hypsugo savii* – Alpenfledermaus. – In: KRAPP, F. (Hrsg.): Handbuch der Säugetiere Europas, Fledertiere II: 911-941.
- HORACEK, I., W. BOGDANOWICZ, & B. DULIC (2004): *Plecotus austriacus* – Graues Langohr. – In: KRAPP, F. (Hrsg.): Handbuch der Säugetiere Europas, Fledertiere II: 1001-1049.
- HORACEK, I. & B. DULIC (2004): *Plecotus auritus* – Braunes Langohr. – In: KRAPP, F. (Hrsg.): Handbuch der Säugetiere Europas, Fledertiere II: 953-999.
- KIEFER, A. & M. VEITH (2001): A new species of long-eared bat from Europe (Chiroptera: Vespertilionidae). – Myotis 39: 5-16.
- KIEFER, A. (2004): Gattung *Plecotus* – Langohrfledermäuse. – In: KRAPP, F. (Hrsg.): Handbuch der Säugetiere Europas, Fledertiere II: 943-952.
- KIEFER, A. & M. MUCEDDA (2004): *Plecotus sardus* – Sardisches Langohr. – In: KRAPP, F. (Hrsg.): Handbuch der Säugetiere Europas, Fledertiere II: 1067-1070.
- KIEFER, A. & O. VON HELVERSEN (2004): *Plecotus macrobullaris* – Alpenlangohr. – In: KRAPP, F. (Hrsg.): Handbuch der Säugetiere Europas, Fledertiere II: 1051-1058.
- KIEFER, A. & O. VON HELVERSEN (2004): *Plecotus kolombatovici* – Balkanlangohr. – In: KRAPP, F. (Hrsg.): Handbuch der Säugetiere Europas, Fledertiere II: 1059-1066.
- KÖNIG, C. (1959): Äußere Merkmale zur Bestimmung der lebenden Fledermäuse Europas. – Säugetierkundliche Mitteilungen 7 (3): 101-110.
- MAYER, F. & O. VON HELVERSEN (2001): Sympatric distribution of two cryptic bat species across Europe. – Biological Journal of the Linnean Society 74: 365-374.
- MAYER, F. & O. VON HELVERSEN (2001): Cryptic diversity in European bats. – Proc. R. Soc. Lond. B 268: 1825-1832.
- MENU, H. & J.-B. POPELARD (1987): Utilisation de caractères dentaires pour la détermination des Vespertilionines de l'Ouest Européen. – Le Rhinolophe 4: 1-88.
- MITCHELL-JONES, A.J., G. AMORI, W. BOGDANOWICZ, B. KRISTUFEK, P.J.H. REIJNDERS, F. SPITZENBERGER, M. STUBBE, J.B.M. THISSEN, V. VOHRALIK & J. ZIMA (1999): The atlas of European mammals, 484 pp.; Poyser Natural History.
- MUCEDDA, M., A. KIEFER, E. PIDINCHEDDA & M. VEITH (2002): A new species of long-eared bat (Chiroptera, Vespertilionidae) from Sardinia (Italy). – Acta Chiropterologica 4 (2): 121-135.

- RUPRECHT, A.L. (1981): Variability of Daubenton's bat and distribution of the *nathalinae* morphotype in Poland. – Acta Theriologica 26: 349-357.
- RUPRECHT, A.L. (1990): Zur Variabilität der Breitflügelfledermäuse und zum Problem um *Eptesicus sodalis* in Polen. – Nyctalus (N.F.) 3 (2): 129-143.
- SCHOBER, W. & E. GRIMMBERGER (1998): Die Fledermäuse Europas. – Kosmos-Naturführer, 265 pp.; Stuttgart.
- SPITZENBERGER, F. (1994): The genus *Eptesicus* (Mammalia, Chiroptera) in southern Anatolia. – Folia Zoologica 43 (4): 437-454.
- SPITZENBERGER, F., J. PIALEK & E. HARING (2001): Systematics of the genus *Plecotus* (Mammalia, Vespertilionidae) in Austria based on morphometric and molecular investigations. – Folia Zool. 50 (3): 161-172.
- SPITZENBERGER, F., E. HARING & N. TVRTKOVIC (2002): *Plecotus microdontus* (Mammalia, Vespertilionidae), a new bat species from Austria. – Nat. Croat. 11 (1): 1-18.
- SPITZENBERGER, F., P. STRELKOV & E. HARING (2003): Morphology and mitochondrial DNA sequences show that *Plecotus alpinus* KIEFER & VEITH, 2002 and *Plecotus microdontus* SPITZENBERGER, 2002 are synonyms of *Plecotus macrobullaris* KUZJAKIN, 1965. – Nat. Croat. 12 (2): 39-53.
- TAAKE, K.-H. (1997): Artbestimmung weiblicher Bartfledermäuse (*Myotis mystacinus/brandti*). – Nyctalus (N.F.), 6 (3): 318.
- TOPAL, G. & M. RUEDI (2001): *Myotis blythii*, Kleines Mausohr. – In Krapp, F. (Hrsg.) (2001): Handbuch der Säugetiere Europas, Band 4: Fledertiere I: 209-255; Aula Verlag.
- TUPINIER, Y. (1977): Description d'une chauve-souris nouvelle: *Myotis nathalinae* nov. sp. – Mammalia 41 (3): 327-340.
- VON HELVERSEN, O. (1989): Bestimmungsschlüssel für die europäischen Fledermäuse nach äußeren Merkmalen. – Myotis 27: 41-60.
- VON HELVERSEN, O. (1998): *Eptesicus bottae* (Mammalia, Chiroptera) auf der Insel Rhodos. – Bonner Zoologische Beiträge 48 (2): 113-121.
- VON HELVERSEN, O., K.-G. HELLER, F. MAYER, A. NEMETH, M. VOLLETH & P. GOMBKÖTÖ (2001): Cryptic mammalian species: a new species of whiskered bat (*Myotis alcathoe* n.sp.) in Europe. – Naturwissenschaften 88: 217-223.
- VON HELVERSEN, O. & M. HOLDERIED (2003): Zur Unterscheidung von Zwergfledermaus (*Pipistrellus pipistrellus*) und Mückenfledermaus (*Pipistrellus mediterraneus/pygmaeus*) im Feld. – Nyctalus (N.F.) 8 (5): 420-426.
- VON HELVERSEN, O. (2004): *Myotis alcathoe* – Nymphenfledermaus. – In: Krapp, F. (Hrsg.): Handbuch der Säugetiere Europas, Fledertiere II: 1159-1167.

APPENDIX 1: LIST OF SCIENTIFIC AND ENGLISH NAMES OF THE EUROPEAN BAT SPECIES

Only the synonyms that have been debated in the last few years (especially of the newly described species) are listed. Reference citations for the newly described species are given in the reference list.

Rhinolophus ferrumequinum (Schreber, 1774) – greater horseshoe bat

Rhinolophus hipposideros (Bechstein, 1800) – lesser horseshoe bat

Rhinolophus euryale Blasius, 1853 – Mediterranean horseshoe bat

Rhinolophus blasii Peters, 1866 – Blasius' horseshoe bat

Rhinolophus mehelyi Matschie, 1901 – Mehely's horseshoe bat

Myotis daubentonii (Kuhl, 1817) – Daubenton's bat

Synonym: *Myotis nathalinae* Tupinier, 1977

Myotis capaccinii (Bonaparte, 1837) – long-fingered bat

Myotis dasycneme (Boie, 1825) – pond bat

Myotis brandtii (Eversmann, 1845) – Brandt's bat

Myotis mystacinus (Kuhl, 1817) – whiskered bat

Myotis aurascens Kusjakin, 1935 – steppe whiskered bat, Eastern whiskered bat

Myotis alcathoe von Helversen & Heller, 2001 – Alcathoe's bat, nymph bat

Myotis emarginatus (Geoffroy, 1806) – Geoffroy's bat

Myotis nattereri (Kuhl, 1817) – Natterer's bat

Myotis bechsteinii (Kuhl, 1817) – Bechstein's bat

Myotis myotis (Borkhausen, 1797) – greater mouse-eared bat

Myotis blythii (Tomes, 1857) – lesser mouse-eared bat

Myotis punicus (Felten, 1977) – Maghrebian mouse-eared bat

Nyctalus noctula (Schreber, 1774) - noctule

Nyctalus leisleri (Kuhl, 1817) – Leisler's noctule

Nyctalus lasiopterus (Schreber, 1780) – greater noctule

Eptesicus serotinus (Schreber, 1774) – serotine

Eptesicus nilssonii (Keyserling & Blasius, 1839) – northern bat

Eptesicus bottae (Peters, 1869) – Botta's serotine

Vespertilio murinus Linnaeus, 1758 – parti-coloured bat

Pipistrellus pipistrellus (Schreber, 1774) – common pipistrelle, 45 kHz pipistrelle

Pipistrellus pygmaeus (Leach, 1825) – soprano pipistrelle, midge bat,
55 kHz pipistrelle
Synonym: *Pipistrellus mediterraneus* Cabrera, 1904

Pipistrellus nathusii (Keyserling & Blasius, 1839) – Nathusius' pipistrelle

Pipistrellus kuhlii (Kuhl, 1817) – Kuhl's pipistrelle

Hypsugo savii (Bonaparte, 1837) – Savi's pipistrelle

Plecotus auritus (Linnaeus, 1758) – common long-eared bat, brown long-eared bat

Plecotus austriacus (Fischer, 1829) – grey long-eared bat

Plecotus macrobullaris (Kusjakin, 1965) – Alpine long-eared bat
Synonyms: *Plecotus alpinus* Kiefer & Veith, 2002
Plecotus microdontus Spitzenberger, 2002

Plecotus kolombatovici (Dulic, 1980) – Kolombatovic's long-eared bat,
Balkan long-eared bat

Plecotus sardus Mucedda & Kiefer, 2002 – Sardinian long-eared bat

Barbastella barbastellus (Schreber, 1774) - barbastelle

Miniopterus schreibersii (Kuhl, 1817) – Schreiber's bat, bent-winged bat

Tadarida teniotis (Rafinesque, 1814) – European free-tailed bat